

The Columbia Lighthouse Project

The Columbia Suicide Severity Rating Scale (C-SSRS)

Supporting Evidence

Last Revised
5-26-2023

Contents

- THE COLUMBIA SUICIDE SEVERITY RATING SCALE (C-SSRS): PSYCHOMETRIC EVIDENCE..... 3**
- TABLE 1: STUDIES SUPPORTING SPECIFIC PSYCHOMETRIC PROPERTIES 3**
- TABLE 2: PSYCHOMETRIC PROPERTIES OF THE C-SSRS IDEATION AND BEHAVIOR WITH COEFFICIENTS 4**
- REFERENCES FOR PSYCHOMETRIC EVIDENCE (TABLES 1&2) 10**
- SCORING AND DATA ANALYSIS GUIDES 12**
- LINGUISTIC AND PSYCHOMETRIC VALIDATION OF THE C-SSRS TRANSLATIONS 12**
- COLUMBIA SUICIDE SEVERITY RATING SCALE VERSIONS 13**
 - C-SSRS Clinical Practice Screener 13**
 - C-SSRS Self-Report 13**
- THE COLUMBIA SUICIDE SEVERITY RATING SCALE (C-SSRS): DIAGNOSTIC AND TREATMENT-MONITORING EFFECTIVENESS 15**
- TABLE 3: 15**
- C-SSRS TRAINING AND QUALITY IMPROVEMENT PROGRAM EVALUATIONS..... 15**
 - Systematic Reviews and Meta-Analyses 16**
- C-SSRS REPRESENTATIVE PUBLICATIONS: DEMOGRAPHIC AND CLINICAL POPULATIONS, SETTINGS, TREATMENT EFFICACY AND ASSESSMENT GUIDELINES..... 16**
 - Pediatric Populations by Age Group 17**
 - Studies with Adolescents and Young Adults..... 20**
- DEMOGRAPHIC GROUPS..... 21**
- SCHOOL PROTOCOLS..... 21**
- MEDICAL SPECIALTIES 21**
 - Neurology..... 21**
 - Oncology..... 21**
 - Psoriasis 22**
- PSYCHIATRIC CONDITIONS..... 22**
 - Alzheimer’s..... 22**
 - ADHD 22**
 - Alcohol 22**
 - Autism 22**
 - Bipolar Depression..... 22**
 - Complicated Grief..... 23**
 - Generalized Anxiety Disorder (GAD)..... 23**
 - Perinatal/Postpartum Depression 23**
 - Psychosis/Schizophrenia 23**
 - PTSD 23**
 - Sleep..... 23**

HEALTHCARE SYSTEMS	24
OUTPATIENT SETTINGS.....	24
Outpatient Psychiatry	24
Integrated Primary Care	24
Veterans	24
Forensic Settings	26
Juvenile Justice	26
EMERGENCY DEPARTMENTS	26
IN-PATIENT PSYCHIATRIC SETTINGS.....	27
MOBILE CRISIS	27
TELEMEDICINE	27
TREATMENT EFFICACY FOR SUICIDAL OUTCOMES	27
REGULATORY GUIDELINES FOR TREATMENT & ASSESSMENT OF SUICIDAL OUTCOMES.....	28
REVIEWS OF SUICIDE RISK ASSESSMENT TOOLS.....	28
CROSS-CULTURAL SETTINGS	28

The Columbia Suicide Severity Rating Scale (C-SSRS): Psychometric Evidence

Table 1: Studies Supporting Specific Psychometric Properties

Psychometric Property		Studies
Clinical Utility	Predictive and/or Incremental Validity	Brent et al., 2009 [^] ; Posner et al., 2011 ^{*^} ; Mundt et al., 2013 [*] ; Arias et al. 2013 [*] ; Greist et al. 2014 [*] ; Jang et al., 2014 [*] ; Brown et al., 2015 [*] ; Gipson et al., 2015 [^] ; Horwitz et al., 2015 [^] ; Arias et al., 2016 [*] ; Madan et al. 2016 [*] ; Conway et al. 2016 [^] ; Hill et al 2017 [^] ; Lindh et al. 2018; Matarazzo et al. 2018 [*] ; Park et al. 2019 ^{*^} ; King et al. 2019 [^] ; Bjureberg et al 2021 [*]
	Sensitivity to Change	Posner et al., 2011 ^{*^} ; Ionescu et al., 2016 [*] ; Lindh et al, 2019 [*]
	Sensitivity and Specificity	Posner et al., 2011 ^{*^} ; Mundt et al., 2013 [*] ; Viguera et al. 2015 [*] ; Madan et al. 2016 [*] ; Lindh et al, 2019 [*] ; Bjureberg et al 2021 [*]
	Positive and Negative Predictive Value (PPV & NPV)	Mundt et al 2013 [*] ; Viguera et al 2015 [*] ; Park et al 2019 [*] ; Bjureberg et al 2021
Reliability (internal consistency)		Posner et al., 2011 ^{*^} ; Kilincaslan et al. 2018 [^] ; Pai et al. 2015 [*] ; Madan et al. 2016 [*] ; Franks et al 2020 [*]
Reliability (inter-rater; multi-method agreement; test-retest)		Kerr et al., 2013 [^] ; Brent et al., 2009 [^] ; Kilincaslan et al. 2018 [^] ; Hesdorffer et al., 2013 [*] ; Arias et al., 2013 [*] ; Brown et al. 2015 [*] ; Tabares et al. 2020 [*] ; Campos et al 2021
Internal Structure (Factor Analysis)		Al-Halabi et al., 2016b [*] ; Madan et al. 2016 [*] ; Tabares et al 2020 [*] ; Franks et al 2020 [*]
Convergent Validity & Accuracy		Posner et al., 2011 ^{*^} ; Kerr et al., 2013 [^] ; Kilincaslan et al. 2018 [^] ; Pai et al. 2015 [*] ; Youngstrom et al. 2015 [*] ; Brown et al., 2015 [*] ; Madan et al.2016 [*] ; Flamarique et al 2016 [^] (concurrent – Campos et al 2021)
Divergent & Discriminant Validity		Posner et al., 2011 ^{*^} ; Kerr et al., 2013 [^] ; Kilincaslan et al. 2018 [^]
Cross-Cultural Validation		Danish (Conway et al. 2016 [^]); Turkish (Kilincaslan et al. 2018 [^]); Korean (Pai et al. 2015 [*]); Spanish (Al-Halabi et al ., 2016ab [*]); Lebanese

* studies include adult samples; ^ studies include pediatric samples

Table 2: Psychometric Properties of the C-SSRS Ideation and Behavior with Coefficients

Predictive Validity - Suicidal Ideation			
	Predictor	Criterion	Coefficients
Bjureberg et al 2021 Adult psych emergency department (N= 18684)	<i>Ideation severity (1-5) past month</i>	Death by suicide	Within 7 days: OR =1.6 (1.2–2.1) <u>Within one month:</u> adjusted OR 1.5, 95% CI 1.2-1.8 Within one year: adjusted OR 1.3, 95% CI 1.1–1.4
	<i>Ideation severity (0-2 vs 3-5) past month</i>	Death by suicide	Within 7 days: Adj OR= 4.7 (1.5–14.8) Within one month: Adj OR= 4.0 (1.9–8.6) Within one year: Adj OR=2.4 (1.5–3.6)
Greist et al. 2014 eCSSRS, drug trials 6,760 patients with psychiatric disorders and 2,077 nonpsychiatric disorder patients	<i>None Reported</i>	Actual, interrupted or aborted attempts	<u>All patients:</u> 0.8% incidence rate, N=4975 <u>Psychiatric patients:</u> 1.1% incidence rate, N=3184
	<i>Wish to Be Dead</i>	Actual, interrupted or aborted attempts	OR= 6.21, 95% CI = 4.18 – 9.23, p <0.001 OR= 4.99, 95% CI = 3.29 – 7.56, p <0.001
	<i>Non-Specific Active Thoughts</i>	Actual, interrupted or aborted attempts	OR= 6.69, 95% CI = 4.16 – 10.76, p <0.001 OR= 5.53, 95% CI = 3.38-9.04, p <0.001
	<i>Active with any methods (not plan) w/o intent to act</i>	Actual, interrupted or aborted attempts	OR= 11.16, 95% CI = 7.43-16.76, p <0.001 OR= 8.36, 95% CI = 5.44-12.84, p <0.001
	<i>Active with Some Intent to Act, without specific plan</i>	Actual, interrupted or aborted attempts	OR= 19.27, 95% CI = 12.97 – 28.63, p <0.001 OR= 15.24, 95% CI = 10.07-23.09, p <0.001
	<i>Active with specific plan and intent</i>	Actual, interrupted or aborted attempts	OR= 25.53, 95% CI = 16.94 – 38.47, p <0.001 OR= 18.70, 95% CI = 12.16 – 28.76, p <0.001
Posner et al. 2011 (TASA study N=124, ages 12-18)	<i>Baseline worst-point</i>	Attempts	OR=1.45, 95% CI=1.07-1.98, p=0.02
		Actual, interrupted and aborted attempts	OR=1.34, 95% CI=1.05-1.70, p=0.02
	<i>Lifetime severity</i>	Attempts	OR=1.43, 95% CI=0.99-2.05, p=0.05
	<i>Severity 4-5 (any intent to</i>	Attempts	OR=3.26, 95% CI=1.02-10.45, p=0.047

	<i>act)</i>	Actual, interrupted and aborted attempts	OR= 3.26, 95% CI=1.07-7.12, p=0.036
Horwitz et al. 2015 (N=473, ages 15-24)	<i>Ideation severity 1 to 5</i>	Attempt	OR= 1.51, 95% CI= 1.24-1.84, p<0.001
Arias et al. 2016 (N=874, mean age 37)	<i>Current ideation severity 4 or 5 (with intent to die)</i>	Actual attempt or suicide 6 weeks post-ED visit	OR=1.70 95% CI 1.18-2.44, p =.004
		Actual, interrupted, aborted attempts, suicide or preparatory behavior	OR =1.52 95%CI 1.23-1.86 p <. 001
Madan et al. 2016: (N=1,055 adult psych in-patients)	<i>Most severe ideation within 72 hours of hospitalization</i>	Any suicide behavior within 6 months post hospitalization	$r = .165, p < .01, N=275$
		Psychiatric re-hospitalization within 6 months	$r = .125, p < .05, N=275$
Conway et al. 2016: (N=85 adolescents, age < 18, mean age=16.2)	Severity of ideation (1-5)	Any type of suicidal behavior at follow-up	OR= 1.66, 95% CI= 1.13-2.44, p<0.05
	Ideation with intent to act (4 or 5)		OR= 7.76, 95% CI= 1.66-36.23, p<0.01
Lindh et al. 2018 (N=804 Adults, age 18-95 years, median age=33)	Most Severe Ideation	Actual Attempt	OR= 1.2, 95% CI= 0.9-1.4, p=0.06
Matarazzo et al 2018 (n=237, mean age 46.1)	<i>Ideation severity 1 to 5</i>	Actual attempt	OR= 2.93 CI= 0.89 - 11.1, p=.02
		Preparatory behavior	OR= 1.95 CI= 1.14-3.32, p < .01
		Any behavior	OR= 1.84 CI= 1.23-2.75, p < .01
Park et al 2019 (N=1359, age≥10)	<i>Ideation severity 1 to 5</i>	Planned actual attempt	OR= 1.58 CI= 1.36-1.83, p < .01
	<i>Ideation severity 5 (with specific plan and any intent)</i>	Planned actual attempt	OR=5.30 CI 1.17-24.07, p < . 05

King et al 2019 (N=2,104, ages 12-17)	<i>Lifetime ideation severity</i>	Actual attempt or death within 3 months follow up	OR = 1.35 CI 1.03 - 1.76, p = .031
Predictive Validity – SI Intensity			
	Predictor	Criterion	Coefficients
Conway et al. 2016: (N=85 adolescents, age < 18, mean age=16.2)	Ideation intensity total score	Any type of suicidal behavior at follow-up	OR= 1.27, 95% CI= 1.04-1.54, p<0.05
Lindh et al. 2018 (N=804 adults, ages 18-95 years, median age=33)	Total Intensity Score	Actual Attempt	OR= 1.07, 95% CI= 1.03-1.1, p=0.001
	Frequency		OR= 1.2, 95% CI= 1.1-1.4, p=0.002
	Duration		OR= 1.2, 95% CI= 1.03-1.3, p=0.01
	Controllability		OR= 1.1, 95% CI= 1.01-1.3, p=0.03
	Deterrents		OR= 1.1, 95% CI= 1.03-1.3, p=0.02
	Reasons		OR= 1.1, 95% CI= 0.9-1.3, p=0.3
Gipson et al. 2014 (N=178, ages 13-17)	Total Intensity Score	Return Psychiatric Emergency Visit	OR= 1.09, 95% CI= 1.00-1.19, p<0.05
	Duration	Return Psychiatric Emergency Visit	OR= 1.67, 95% CI= 1.16-2.42, p<0.01
		Actual Attempt	OR= 1.80, 95% CI= 1.06-3.04, p<0.05
Predictive Validity - Suicidal Behavior			
	Predictor	Criterion	Coefficients
Bjureberg et al 2021	<i>Suicidal behavior question</i>	Death by suicide	Within 7 days: adjusted OR 6.9, 95% CI 2.1– 22.7 Within one month: adjusted OR 5.1, 95% CI 2.3-11.2) Within one year: adjusted OR 2.8, 95% CI 1.7–4.5

Horwitz et al. 2015: (N=473, ages 15-24)	<i>Attempt</i>	Attempt	OR=4.80, 95% CI = 2.23-10.32, p<0.001
	<i>NSSIB item</i>	Attempt	OR=3.12, 95% CI = 1.36-7.19, p<0.01
Gipson et al. 2014 (N=178, ages 13-17)	<i>NSSIB item</i>	Return ER visit	OR = 1.52; 95% CI, 1.08-2.12, p<.05
		Attempt	$\chi^2 = 4.131$, df = 1, p = 0.04
Conway et al. 2016 (N=85, age < 18, mean age=16.2)	<i>Attempts</i>	Re-attempt [short-term]	OR= 11.50, 95% CI= 1.66-79.65, p<0.05
Greist et al. 2014	<i>Attempt</i>	Actual, interrupted or aborted attempts	OR=4.57, 95% CI = 3.6-5.7, p<0.001
	<i>Interrupted Attempt</i>	Actual, interrupted or aborted attempts	OR=5.55, 95% CI = 4.4-7.0, p<0.001
	<i>Aborted Attempt</i>	Actual, interrupted or aborted attempts	OR=5.09, 95% CI = 4.1-6.4, p<0.001
	<i>Preparatory behavior</i>	Actual, interrupted or aborted attempts	OR=5.69, 95% CI = 4.3-7.5, p<0.001

Incremental Validity and Accuracy

Brent et al., (2009): Treatment resistant, depressed adolescent suicide attempters (N=334, ages 12-18)	<ul style="list-style-type: none"> Higher rates of suicidal (20.8% vs. 8.8%, chi squared= 9.18, df=1, p<0.002) and non-suicidal self-injury (17.6% vs. 2.2%, chi squared= 23.47, df=1, p<0.001) detected with systematic monitoring
Horwitz et al. (2015): Young adult psychiatric emergency patients (N=473, ages 15-24)	<ul style="list-style-type: none"> Suicidal ideation added incremental validity to the prediction of future suicide attempts beyond the past suicide attempt, $\chi^2 (1) = 7.54$, p= .006
Brown et al. (2015): psychiatric ER patients (N=250)	<ul style="list-style-type: none"> 18% (n=23) of patients with a suicide attempt in the past week misclassified or missed by clinical assessment. Agreement with clinical assessment for suicide attempts (K=0.76, p=<.001) Agreement with clinical assessment of non-suicidal self-injurious behavior (K=0.72, p=<.001)
Arias et al. (2013): 497 ER adult patients with suicidal thoughts or attempt(s)	<ul style="list-style-type: none"> 41% increase in the detection of suicide attempts compared to chart reviews (59% vs. 18%, difference of 41%, 95% CI= 28-55, p<0.001)

Reliability - Suicidal Ideation (inter-rater and multi-method agreement)		
Study	Ideation Type	Coefficients
Brent et al. (2009) (N=334, ages 12-18)	<i>suicidal ideation ranging from 0 to 5 (from no ideation to suicidal ideation with intent and a clear plan) monitored weekly</i>	ICC = .09, p< 0.001
Kilincaslan et al. (2018) (N=213, ages 12-18)	<i>Inter-rater reliability for the <u>most severe ideation scores</u> in the last month and lifetime were good</i>	Lifetime κ = 0.92 Recent κ = 0.88
Youngstrom et al. (2015)	<i>Accuracy calibrated against “missing gold standard” latent class-derived ideation and behavior categories</i>	$\kappa > 0.7$
Hesdorffer et al. (2013)	<i>Agreement between the MINI, C-SSRS and eC-SSRS for lifetime <u>suicidal ideation</u></i>	κ = 0.80, 95% CI = 0.72-0.89
Gwaltney et al. (2017) (N=86, ages >18)	<i>Equivalence analyses/multi-method agreement between IVR (interactive voice response) and tablet text-based eC-SSRS for <u>most severe lifetime ideation</u></i>	Correlation: 0.87, p<0.001 ICC: κ = 0.89, p<0.001
	<i>Equivalence analyses between IVR (interactive voice response) and tablet text-based eC-SSRS for <u>most severe ideation in past 6 months</u></i>	Correlation: 0.69, p<0.001 ICC: κ = 0.79, p<0.001
Reliability - Suicidal Behavior		
Gwaltney et al. (2017) (N=86, ages >18)	<i>Equivalence analyses/multi-method agreement between IVR (interactive voice response) and tablet text-based eC-SSRS for lifetime Actual attempts</i>	κ = 0.81, p<0.001
	<i>Number of lifetime actual attempts</i>	κ = 0.81, p<0.001
	<i>Actual attempts (recent-last 2 yrs)</i>	κ = 0.73, p<0.001

	<i>Interrupted attempts (lifetime)</i>	$\kappa = 0.78, p < 0.001$
	<i>Interrupted attempts (recent-last 2 yrs)</i>	$\kappa = 0.762, p < 0.001$
	<i>Aborted attempts (lifetime)</i>	$\kappa = 0.54, p < 0.001$
	<i>Aborted attempts (recent-last 2 yrs)</i>	$\kappa = 0.74, p < 0.001$
	<i>Preparatory behaviors (lifetime)</i>	$\kappa = 0.77, p < 0.001$
	<i>Preparatory behaviors (recent-last 2 yrs)</i>	$\kappa = 0.89, p < 0.001$
	<i>Non-suicidal, self-injurious behavior</i>	$\kappa = 0.73, p < 0.001$
Brent et al. (2009) (N=334, ages 12-18)	<i>Inter-rater reliability for a rating of <u>suicidal behavior</u>, ranging from 0 to 5 (no behavior to multiple attempts during the assessment period) using the Columbia Classification Algorithm of Suicide Assessment</i>	100% agreement
Kerr et al. (2014a,b) (N=155, ages 13-17)	<i>Inter-rater agreement for distinction among <u>actual, aborted, interrupted attempts, preparatory acts and any other act</u></i>	$\kappa = 0.88; \kappa = .91$
Brown et al. (2015)	<i>Agreement with clinical assessment for <u>attempts</u></i>	$\kappa = 0.76, P < .001$
	<i>Agreement with clinical assessment for <u>non-suicidal self-injurious behavior</u></i>	$\kappa = 0.72, P < .001$
Youngstrom et al. (2015)	<i>Accuracy of <u>attempt</u>: calibrated against latent class-derived categories</i>	$\kappa > 0.8$
Hesdorffer et al. (2013)	<i>Agreement between the MINI, C-SSRS and eC-SSRS for lifetime <u>suicidal behavior</u></i>	$\kappa = 0.67, 95\% \text{ CI} = 0.53-0.80$

References for Psychometric Evidence (Tables 1&2)

1. Al-Halabí, S., Sáiz, P. A., Burón, P., Garrido, M., Benabarre, A., Jiménez, E., ... & Muñiz, J. (2016a). Validación de la versión en español de la Columbia-Suicide Severity Rating Scale (Escala Columbia para Evaluar el Riesgo de Suicidio). *Revista de Psiquiatría y Salud Mental*.
2. Al-Halabi, S., Fernández-Peláez, AD, Burón, P., Riesco, E., Rodríguez-Revuelta, J. Posner, K. Oquendo, M., García-Portilla, MP, Saiz, P. and Bobes, J (September, 2016b). In Search of the Internal Structure of the Columbia Suicide Severity Rating Scale (C-SSRS): A Confirmatory Factor Analysis Approach. *16th European Symposium on Suicide Suicidal Behavior, Oviedo, Spain*.
3. Arias, S. A., Miller, I., Camargo Jr, C. A., Sullivan, A. F., Goldstein, A. B., Allen, M. H., ... & Boudreaux, E. D. (2016). Factors Associated with Suicide Outcomes 12 Months After Screening Positive for Suicide Risk in the Emergency Department. *Psychiatric Services, 67 (2):* 206-213.
4. Arias SA, Zhang Z, Hillerns C, Sullivan AF, Boudreaux ED, Miller I, Camargo CA (2014). Using Structured Telephone Follow-up Assessments to Improve Suicide-Related Adverse Event Detection. *Suicide and Life-Threatening Behavior*44(5): 537-47.
5. Brent D, Emslie G, Clarke G, Rosenbaum Asarnow J, Spirito A, Ritz L, Vitiello B, Iyengar S, Birmaher B, Ryan N, Zelazny J, Onorato M, Kennard B, Mayes T, DeBar L, McCracken J, Strober M, Suddath R, Leonard H, Porta G, Keller M (2009) Predictors of Spontaneous and Systematically Assessed Suicidal Adverse Events in the Treatment of SSRI Resistant Depression in Adolescents (TORDIA) Study. *American Journal of Psychiatry* 166(4): 418-426.
6. Bjureberg, J., Dahlin, M., Carlborg, A., Edberg, H., Haglund, A., & Runeson, B. (2021). Columbia-Suicide Severity Rating Scale Screen Version: initial screening for suicide risk in a psychiatric emergency department. *Psychological medicine, 1-9*.
7. Brown GK, Currier GW, Jager-Hyman S, et al. (2015) Detection and classification of suicidal behavior and nonsuicidal self-injury behavior in emergency departments. *J Clin Psychiatry* 76(10):1397–1403.
8. Campos, A. I., Van Velzen, L. S., Veltman, D. J., Pozzi, E., Ambrogi, S., Ballard, E. D., ... & Rentería, M. E. (2021). Concurrent validity and reliability of suicide risk assessment instruments: A meta analysis of 20 instruments across 27 international cohorts. *medRxiv*.
9. Conway, P. M., Erlangsen, A., Teasdale, T. W., Jakobsen, I. S., & Larsen, K. J. (2016). Predictive Validity of the Columbia-Suicide Severity Rating Scale for Short-Term Suicidal Behavior: a Danish study of adolescents at a high risk of suicide. *Archives of suicide research, 1-15*.
10. Flamarique, I., Santosh, P., Zuddas, A. *et al*. Development and psychometric properties of the Suicidality: Treatment Occurring in Paediatrics (STOP) Suicidality Assessment Scale (STOP-SAS) in children and adolescents. *BMC Pediatr* **16**, 213 (2016).
11. Gipson PY, Agarwala P, Opperman KJ, Horwitz A, King CA, (2015). Columbia-Suicide Severity Rating Scale: Predictive Validity with Adolescent Psychiatric Emergency Patients. *Pediatric Emergency Care* 31(2): 88-94.
12. Greist, J. H., Mundt, J. C., Gwaltney, C. J., Jefferson, J. W., & Posner, K. (2014). Predictive Value of Baseline Electronic Columbia–Suicide Severity Rating Scale (eC–SSRS) Assessments for Identifying Risk of Prospective Reports of Suicidal Behavior During Research Participation. *Innovations in Clinical Neuroscience, 11(9-10), 23–31*.
13. Gwaltney, C., Mundt, J. C., Greist, J. H., Paty, J., & Tiplady, B. (2017). Interactive Voice Response and Text-based Self-report Versions of the Electronic Columbia-Suicide Severity Rating Scale Are Equivalent. *Innovations in clinical neuroscience, 14(3-4), 17*.
14. Harvey, P. D., Posner, K., Rajeevan, N., Yershova, K. V., Aslan, M., & Concato, J. (2018). Suicidal ideation and behavior in US veterans with schizophrenia or bipolar disorder. *Journal of Psychiatric Research*.
15. Hill, R. M., Hatkevich, C. E., Kazimi, I., & Sharp, C. (2017). The Columbia-Suicide Severity Rating Scale: Associations between interrupted, aborted, and actual suicide attempts among adolescent inpatients. *Psychiatry research, 255, 338-340*.
16. Hesdorffer DC, French JA, Posner K, DiVentura B, Pollard J, Sperling MR, Harden CL, Krauss GL, Kanner AM (2013). Suicidal Ideation and behavior screening in intractable focal epilepsy eligible for drug trials. *Epilepsia* 54(5): 879-87.

17. Horwitz, A. G., Czyz, E. K., & King, C. A. (2015). Predicting future suicide attempts among adolescent and emerging adult psychiatric emergency patients. *Journal of Clinical Child & Adolescent Psychology*, 44(5), 751-761.
18. Ionescu, D. F., Swee, M. B., Pavone, K. J., Taylor, N., Akeju, O., Baer, L., ... & Brown, E. N. (2016). Rapid and sustained reductions in current suicidal ideation following repeated doses of intravenous ketamine: secondary analysis of an open-label study. *The Journal of clinical psychiatry*.
19. Jang HA, Park EH, Jon DI, Park HJ, Hong HJ, Jung MH, et al. 2014. Validation of the Columbia Suicide Severity Rating Scale in Depression Patient. *Kor J Clin Psychol*. 33:183–188.
20. Kerr DCR, Gibson B, Leve LD, DeGarmo DS (2013). Young Adult Follow-up of Adolescent Girls in Juvenile Justice Using the Columbia Suicide Severity Rating Scale. *Suicide and Life Threatening Behavior* 44(2): 113-129.
21. Kerr, D. C., DeGarmo, D. S., Leve, L. D., & Chamberlain, P. (2014). Juvenile justice girls' depressive symptoms and suicidal ideation 9 years after multidimensional treatment foster care. *Journal of consulting and clinical psychology*, 82(4), 684.
22. King, C. A., Grupp-Phelan, J., Brent, D., Dean, J. M., Webb, M., Bridge, J. A., ... & Rea, M. (2019). Predicting 3-month risk for adolescent suicide attempts among pediatric emergency department patients. *Journal of child psychology and psychiatry*.
23. Kilincaslan, A., Gunes, A., Eskin, M., & Madan, A. (2018). Linguistic adaptation and psychometric properties of the Columbia-Suicide Severity Rating Scale among a heterogeneous sample of adolescents in Turkey. *The International Journal of Psychiatry in Medicine*, 0091217418791454.
24. Legarreta, M., Graham, J., North, L., Bueler, C. E., McGlade, E., & Yurgelun-Todd, D. (2015). DSM-5 posttraumatic stress disorder symptoms associated with suicide behaviors in veterans. *Psychological trauma: theory, research, practice, and policy*, 7(3), 277.
25. Lindh ÅU, Dahlin M, Beckman K, Strömsten L, Jokinen J, Wiktorsson S, Renberg ES, Waern M, Runeson B. A (2019) Comparison of Suicide Risk Scales in Predicting Repeat Suicide Attempt and Suicide: A Clinical Cohort Study. *J Clin Psychiatry*
26. Madan, A., Frueh, B. C., Allen, J. G., Ellis, T. E., Rufino, K. A., Oldham, J. M., & Fowler, J. C. (2016). Psychometric reevaluation of the Columbia-Suicide Severity Rating Scale: findings from a prospective, inpatient cohort of severely mentally ill adults. *Journal of clinical psychiatry*, 77(7), e867-e873.
27. Matarazzo, B. B., Brown, G. K., Stanley, B., Forster, J. E., Billera, M., Currier, G. W., ... & Brenner, L. A. (2018). Predictive Validity of the Columbia-Suicide Severity Rating Scale among a Cohort of At-risk Veterans. *Suicide and Life-Threatening Behavior*.
28. Mundt JC, Greist JH, Jefferson JW, Federica M, Mann JJ, Posner K (2013). Prediction of Suicidal Behavior in Clinical research by Lifetime Suicidal ideation and Behavior Ascertained by the Electronic Columbia-Suicide Severity Rating Scale. *Journal of Clinical Psychiatry* 74(9):887-93.
29. Posner K, Brown GK, Stanley B, Brent DA, Yershova KV, Oquendo MA, Currier GW, Melvin GA, Greenhill L, Shen S, Mann JJ (2011). The Columbia-Suicide Severity Rating Scale: Initial Validity and Internal Consistency Findings from Three Multisite Studies with Adolescents and Adults. *American Journal of Psychiatry* 168(12): 1266-77.
30. Park, C. H. K., Kim, H., Kim, B., Kim, E. Y., Lee, H. J., Kim, D., & Ahn, Y. M. (2019). Predicting Planned Suicide Attempts With the Columbia-Suicide Severity Rating Scale: A Subanalysis of the 2013 Korea National Suicide Survey on Emergency Department Visitors. *The Journal of nervous and mental disease*, 207(2), 59-68.
31. Park, E. H., Hong, N., Jon, D. I., Hong, H. J., & Jung, M. H. (2017). Past suicidal ideation as an independent risk factor for suicide behaviours in patients with depression. *International journal of psychiatry in clinical practice*, 21(1), 24-28.
32. Prakash, A., Lobo, E., Kratochvil, C. J., Tamura, R. N., Pangallo, B. A., Bullok, K. E., ... & March, J. S. (2012). An open-label safety and pharmacokinetics study of duloxetine in pediatric patients with major depression. *Journal of child and adolescent psychopharmacology*, 22(1), 48-55.
33. Tabares, J. V., Butner, J. E., Bryan, C. J., & A. Harris, J. (2020). Mokken Scale Analysis of Lifetime Responses on the Columbia Suicide Severity Rating Scale's Severity of Ideation Subscale. *Assessment*, 1073191120913626.

34. Viguera, A. C., Milano, N., Laurel, R., Thompson, N. R., Griffith, S. D., Baldessarini, R. J., & Katzan, I. L. (2015). Comparison of electronic screening for suicidal risk with the Patient Health Questionnaire Item 9 and the Columbia Suicide Severity Rating Scale in an outpatient psychiatric clinic. *Psychosomatics*, 56(5), 460-469.
35. Youngstrom, E. A., Hameed, A., Mitchell, M. A., Van Meter, A. R., Freeman, A. J., Algorta, G. P., ... & Meyer, R. E. (2015). Direct comparison of the psychometric properties of multiple interview and patient-rated assessments of suicidal ideation and behavior in an adult psychiatric inpatient sample. *The Journal of clinical psychiatry*, 76(12), 1676-1682.

Other predictive validity studies:

Wilimitis, D., Turer, R. W., Ripperger, M., McCoy, A. B., Sperry, S. H., Fielstein, E. M., ... & Walsh, C. G. (2022). Integration of face-to-face screening with real-time machine learning to predict risk of suicide among adults. *JAMA network open*, 5(5), e2212095-e2212095.

Scoring and Data Analysis Guides

- Nilsson, M. E., Suryawanshi, S., Gassmann-Mayer, C., Dubrava, S., McSorley, P., & Jiang, K. (2013). Columbia–suicide severity rating scale scoring and data analysis guide. *CSSRS Scoring Version, 2*, 1-13.
- Crowe, B., Xia, A., Nilsson, M., Shahin, S., Wang, W., & Jiang, Q. (2015). The program safety analysis plan: An implementation guide. *Quantitative Evaluation of Safety in Drug Development: Design, Analysis, and Reporting*, 55-68.

Resource Utilization

Syndergaard, S., Borger, J., Klenzak, S., Grello, A., & Adams, A. (2022). Implementation of Columbia Suicide Severity Rating Scale (C-SSRS) as a universal suicide risk screening tool in a high-volume emergency department. *Archives of suicide research*, 1-11.

Linguistic and Psychometric Validation of the C-SSRS Translations

- Gatalup, G., Fernander, N., Fuller, D.S. and Posner, K (2013). Translation of the Columbia Suicide Severity Rating Scale for Use in 33 Countries. ISCTM 9th Annual Scientific Meeting, Washington D.C.
- Al-Halabí, S., Sáiz, P. A., Burón, P., Garrido, M., Benabarre, A., Jiménez, E., ... & Muñiz, J. (2016). Validación de la versión en español de la Columbia-Suicide Severity Rating Scale (Escala Columbia para Evaluar el Riesgo de Suicidio). *Revista de Psiquiatría y Salud Mental*, 9(3), 134-142.
- Al-Halabi, S., Fernández-Peláez, AD, Burón, P., Riesco, E., Rodríguez-Revuelta, J. Posner, K. Oquendo, M., García-Portilla, MP, Saiz, P. and Bobes, J (September, 2016). In Search of the Internal Structure of the Columbia Suicide Severity Rating Scale (C-SSRS): A Confirmatory Factor Analysis Approach. *16th European Symposium on Suicide Suicidal Behavior*, Oviedo, Spain. **[Spanish]**
- Conway, P. M., Erlangsen, A., Teasdale, T. W., Jakobsen, I. S., & Larsen, K. J. (2016). Predictive Validity of the Columbia-Suicide Severity Rating Scale for Short-Term Suicidal Behavior: a **Danish** study of **adolescents** at a high risk of suicide. *Archives of suicide research*, 1-15.

- Gunes A, Kilincaslan A, Eskin M (2015). Psychometric Properties of the **Turkish** Version of Columbia-Suicide Severity Rating Scale Among **12-18 year-old** adolescents in Turkey. AACAP 62nd Annual Meeting, San Antonio, TX.
- Pai, D., Woo, J. M., Son, M. H., & Lee, C. (2015). The Reliability and Validity of the Korean Version of Columbia-Suicide Severity Rating Scale in Alcohol Dependent Patients. *Journal of Korean Neuropsychiatric Association*, 54(2), 222-227.
- Serrani Azcurra, D. (2017). Psychometric validation of the Columbia-Suicide Severity rating scale in Spanish-speaking adolescents. *Colombia Médica*, 48(4), 174-182.

Columbia Suicide Severity Rating Scale Versions

(adapted versions)

C-SSRS Clinical Practice Screener

- Bjureberg, J., Dahlin, M., Carlborg, A., Edberg, H., Haglund, A., & Runeson, B. (2021). Columbia-Suicide Severity Rating Scale Screen Version: initial screening for suicide risk in a psychiatric emergency department. *Psychological medicine*, 1-9.
- Katz, I., Barry, C. N., Cooper, S. A., Kaspro, W. J., & Hoff, R. A. (2019). Use of the Columbia-Suicide Severity Rating Scale (C-SSRS) in a large sample of Veterans receiving mental health services in the Veterans Health Administration. *Suicide and Life-Threatening Behavior*.
- Brahmabhatt, K., Kurtz, B. P., Afzal, K. I., Giles, L. L., Kowal, E. D., Johnson, K. P., ... & Workgroup, P. (2019). Suicide risk screening in pediatric hospitals: clinical pathways to address a global health crisis. *Psychosomatics*, 60(1), 1-9.
- Keaton, S. A., Madaj, Z. B., Heilman, P., Smart, L., Grit, J., Gibbons, R., ... & Brundin, L. (2019). An inflammatory profile linked to increased suicide risk. *Journal of affective disorders*, 247, 57-65.
- Imran, J. B., Richmond, R. E., Madni, T. D., Roaten, K., Clark, A. T., Huang, E., ... & Eastman, A. L. (2018). Determining Suicide Risk in Trauma Patients Using a Universal Screening Program. *The journal of trauma and acute care surgery*.
- Henderson, J. L., Cheung, A., Cleverley, K., Chaim, G., Moretti, M. E., de Oliveira, C., ... & Herzog, T. (2017). Integrated collaborative care teams to enhance service delivery to youth with mental health and substance use challenges: protocol for a pragmatic randomised controlled trial. *BMJ*, 7(2).
- Roaten, K., Johnson, C., Genzel, R., Khan, F., & North, C. S. (2017). Development and implementation of a universal suicide risk screening program in a safety-net hospital system. *The Joint Commission Journal on Quality and Patient Safety*.
- Wilson, N. J. (2017). The Columbia-Suicide Severity Rating Scale: Validation for use as a screen for suicide risk in New Zealand prisons and probation settings. *The New Zealand Corrections Journal*, Vol. 5 Issue 2.
- Omolewa, P., & Tribble, K. L. The Impact of C-SSRS (Columbia-Suicidal Severity Rating Scale) Usage on Quality of Care in John George Psychiatric Hospital (San Leandro, CA): a Medical Care Evaluation Study.
- Chang, B. P., & Tan, T. M. (2015). Suicide screening tools and their association with near-term adverse events in the ED. *The American journal of emergency medicine*, 33(11), 1680-1683. [self-report screener]

C-SSRS Self-Report

("paper" non-adaptive version, not eC-SSRS)

- Roaten, K., Johnson, C., Genzel, R., Khan, F., & North, C. S. (2017). Development and implementation of a universal suicide risk screening program in a safety-net hospital system. *The Joint Commission Journal on Quality and Patient Safety*.
- DeVlyder, J. E., Jahn, D. R., Doherty, T., Wilson, C. S., Wilcox, H. C., Schiffman, J., & Hilimire, M. R. (2015). Social and psychological contributions to the co-occurrence of sub-threshold psychotic experiences and suicidal behavior. *Social psychiatry and psychiatric epidemiology*, 1-12. [study adapted self-report]
- Viguera, A. C., Milano, N., Laurel, R., Thompson, N. R., Griffith, S. D., Baldessarini, R. J., & Katzan, I. L. (2015). Comparison of electronic screening for suicidal risk with the Patient Health Questionnaire Item 9 and the Columbia Suicide Severity Rating Scale in an outpatient psychiatric clinic. *Psychosomatics*, 56(5), 460-469.
- Nock, M. K., Ursano, R. J., Heeringa, S. G., Stein, M. B., Jain, S., Raman, R., ... & Gilman, S. E. (2015). Mental disorders, comorbidity, and pre-enlistment suicidal behavior among new soldiers in the US Army: results from the Army Study to Assess Risk and Resilience in Service members (Army STARRS). *Suicide and life-threatening behavior*, 45(5), 588-599.
- Chang, B. P., & Tan, T. M. (2015). Suicide screening tools and their association with near-term adverse events in the ED. *The American journal of emergency medicine*, 33(11), 1680-1683. [self-report screener]
- Katz, I., Barry, C. N., Cooper, S. A., Kasprow, W. J., & Hoff, R. A. (2019). Use of the Columbia-Suicide Severity Rating Scale (C-SSRS) in a large sample of Veterans receiving mental health services in the Veterans Health Administration. *Suicide and Life-Threatening Behavior*. [self-report screener]

The Columbia Suicide Severity Rating Scale (C-SSRS): Diagnostic and Treatment-Monitoring Effectiveness

Table 3:

C-SSRS as an Effective Measure for Diagnosis & Treatment	Veterans Legarreta et al., 2015	<ul style="list-style-type: none"> The association of specific PTSD symptoms with suicidal ideation and behavior suggested individual PTSD symptoms as treatment target for reducing suicidal outcomes.
	Veterans Harvey et al., 2018	<ul style="list-style-type: none"> A lifetime history of suicidal ideation and behavior was higher among the Vets with Bipolar Disorder (82.3%, N=5414) than Schizophrenia (69.9%, N=3942) The highest risk was found for patients with multiple psychiatric comorbidities (OR = 2.61 for ideation; OR = 3.82 for behavior). Clinical factors (e.g., psychiatric comorbidity) contributed more of the variance in the predictive model than demographic factors.
	Medication Treatment Ionescu et al. (2016) Prakash et al. (2012)	<ul style="list-style-type: none"> Ketamine treatment effective for suicidal ideation (SI) in adults SI severity improved <u>independent</u> of acute decrease in depression and SI intensity improved <u>even if SI severity un-remitted</u> Duloxetine was effective in treating suicidal ideation among children ages 7-17 with major depression Distinguished children with improvement and deterioration

C-SSRS Training and Quality Improvement Program Evaluations

- Fesi, J. D., Morrison, S. U. (2018, 04). Guardians At the Gate: Evaluating Suicide Risk Assessment Training in the United States Marine Corps. 2018 American Association of Suicidology 51st Annual Conference, Washington D.C.
- Mirick, R. G., Bridger, J., McCauley, J., & Berkowitz, L. (2016). Continuing Education on Suicide Assessment and Crisis Intervention for Social Workers and Other Mental Health Professionals: A Follow-Up Study. *Journal of Teaching in Social Work*, 36(4), 363-379.
- Mirick, R., McCauley, J., Bridger, J., & Berkowitz, L. (2015). Continuing education on suicide assessment and crisis intervention: what can we learn about the needs of mental health professionals in community practice? *Community mental health journal*, 1-10.
- Soffer et al (2022) Assessing Suicide Risk in a Pediatric Outpatient Behavioral Health System: A Quality Improvement Report. *Pediatr Qual Saf*, 7(3).

- Latif, F., Patel, S., Badolato, G., McKinley, K., Chan-Salcedo, C., Bannerman, R., ... & Robb, A. S. (2020). Improving youth suicide risk screening and assessment in a pediatric hospital setting by using the joint commission guidelines. *Hospital Pediatrics*, 10(10), 884-892.
- Epilepsy: Rani, R. A., Razali, R., Hod, R., Mohamad, K., Rani, S. A. M., Wan Yahya, W. N. N., ... & Tan, H. J. (2014). Suicidal ideation amongst epilepsy patients in a tertiary centre. *Neurology Asia*, 19(2).
- Emerson, B. L., Setzer, E., Blake, E., & Siew, L. (2022). Improving Quality and Efficiency in Pediatric Emergency Department Behavioral Health Care. *Pediatric Quality & Safety*, 7(1).

COVID-19 Studies

- Patella, V., Pelaia, C., Zunno, R., & Pelaia, G. (2022). Biologicals decrease psychological distress, anxiety and depression in severe asthma, despite Covid-19 pandemic. *Respiratory Medicine*, 200, 106916.
- Hill et al (2021). Suicide ideation and attempts in a pediatric emergency department before and during COVID-19. *Pediatrics*, 147.
- Hill et al (2021). Geospatial identification of high youth suicide risk areas via electronic health records: avenues for research and prevention efforts. *Suicide Life Threaten Behav.*, 51:255–62.

Systematic Reviews and Meta-Analyses

- Carter, T., Walker, G. M., Aubeeluck, A., & Manning, J. C. (2019). Assessment tools of immediate risk of self-harm and suicide in children and young people: a scoping review. *Journal of child health care*, 23(2), 178-199.
- Scudder A, Rosin R, Baltich Nelson B, Boudreaux ED, Larkin C. (2022) Suicide Screening Tools for Pediatric Emergency Department Patients: A Systematic Review. *Front Psychiatry*. 2022 Jul 12;13.
- Harris, I. M., Beese, S., & Moore, D. (2019). Predicting future self-harm or suicide in adolescents: a systematic review of risk assessment scales/tools. *BMJ open*, 9(9).
- Riblet, N. B., Matsunaga, S., Lee, Y., Young-Xu, Y., Shiner, B., Schnurr, P. P., ... & Watts, B. V. (2022). Tools to detect risk of death by suicide: a systematic review and meta-analysis. *The Journal of Clinical Psychiatry*, 84(1), 43891.
- Campos, A. I., Van Velzen, L. S., Veltman, D. J., Pozzi, E., Ambroggi, S., Ballard, E. D., ... & Rentería, M. E. (2021). Concurrent validity and reliability of suicide risk assessment instruments: A meta-analysis of 20 instruments across 27 international cohorts. medRxiv.

C-SSRS Representative Publications: Demographic and Clinical Populations, Settings, Treatment Efficacy and Assessment Guidelines

Pediatric Populations by Age Group

Ages 5-11

- Glennon, J., Purper-Ouakil, D., Bakker, M., Zuddas, A., Hoekstra, P., Schulze, U., ... & Coghill, D. (2014). Paediatric European Risperidone Studies (PERS): context, rationale, objectives, strategy, and challenges. *European child & adolescent psychiatry*, 23(12), 1149-1160. [also includes 12-17.5 age group]
- Proskurina, T., Mykhailova, E., Matkovska, T., Reshetovska, N., & Matkovska, A. (2016). Clinical and psychopathological risk factors for the development of recurrent depression in children at puberty. *European Psychiatry*, 33, S145-S146.
- Sheftall, A. H., Vakil, F., Armstrong, S. E., Rausch, J. R., Feng, X., Kerns, K. A., ... & Bridge, J. A. (2021). Clinical risk factors, emotional reactivity/regulation and suicidal ideation in elementary school-aged children. *Journal of psychiatric research*, 138, 360-365.

Ages 6-12

- Buchanan, J., Burke, T., Camacho, K., Yershova, K., Lazzaretto, D., Posner, K. (2013) Preschool Bullying and Victimization as Predictors of Suicidal Ideation in School Age: 6-year Follow-Up of the Preschool Attention Deficit/Hyperactivity Disorder Treatment Study (PATS). *1st Annual Meeting of the International Academy for Suicide Research*, Montreal, Canada.
- Childress, A. C., Wigal, S. B., Brams, M. N., Turnbow, J. M., Pincus, Y., Belden, H. W., & Berry, S. A. (2018). Efficacy and safety of amphetamine extended-release oral suspension in children with attention-deficit/hyperactivity disorder. *Journal of child and adolescent psychopharmacology*, 28(5), 306-313.
- Latif, F., Patel, S., Badolato, G., McKinley, K., Chan-Salcedo, C., Bannerman, R., ... & Robb, A. S. (2020). Improving youth suicide risk screening and assessment in a pediatric hospital setting by using the joint commission guidelines. *Hospital Pediatrics*, 10(10), 884-892

Ages 6-17

- Glennon, J., Purper-Ouakil, D., Bakker, M., Zuddas, A., Hoekstra, P., Schulze, U., ... & PERS Consortium. (2014). Paediatric European Risperidone Studies (PERS): context, rationale, objectives, strategy, and challenges. *European child & adolescent psychiatry*, 23(12), 1149-1160.
- Sangal, R. B., Blumer, J. L., Lankford, D. A., Grinnell, T. A., & Huang, H. (2014). Eszopiclone for insomnia associated with attention-deficit/hyperactivity disorder. *Pediatrics*, 134(4), e1095-e1103.
- Latif, F., Patel, S., Badolato, G., McKinley, K., Chan-Salcedo, C., Bannerman, R., ... & Robb, A. S. (2020). Improving youth suicide risk screening and assessment in a pediatric hospital setting by using the joint commission guidelines. *Hospital Pediatrics*, 10(10), 884-892.

Ages 6-18

- Wigal, S. B., Nordbrock, E., Adjei, A. L., Childress, A., Kupper, R. J., & Greenhill, L. (2015). Efficacy of Methylphenidate Hydrochloride Extended-Release Capsules (Aptensio XR™) in Children and Adolescents with Attention-Deficit/Hyperactivity Disorder: A Phase III, Randomized, Double-Blind Study. *CNS drugs*, 29(4), 331-340.

Ages 7-13

- Weinstein, S. M., Henry, D. B., Katz, A. C., Peters, A. T., & West, A. E. (2015). Treatment moderators of child-and family-focused cognitive-behavioral therapy for pediatric bipolar disorder. *Journal of the American Academy of Child & Adolescent Psychiatry*, 54(2), 116-125.

Ages 7-17

- Strawn, J. R., Prakash, A., Zhang, Q., Pangallo, B. A., Stroud, C. E., Cai, N., & Findling, R. L. (2015). A randomized, placebo-controlled study of duloxetine for the treatment of children and adolescents with generalized anxiety disorder. *Journal of the American Academy of Child & Adolescent Psychiatry*, 54(4), 283-293.
- Emslie, G. J., Prakash, A., Zhang, Q., Pangallo, B. A., Bangs, M. E., & March, J. S. (2014). A double-blind efficacy and safety study of duloxetine fixed doses in children and adolescents with major depressive disorder. *Journal of child and adolescent psychopharmacology*, 24(4), 170-179.
- Prakash, A., Lobo, E., Kratochvil, C. J., Tamura, R. N., Pangallo, B. A., Bullok, K. E., ... & March, J. S. (2012). An open-label safety and pharmacokinetics study of duloxetine in pediatric patients with major depression. *Journal of child and adolescent psychopharmacology*, 22(1), 48-55.

Ages 7-18 (for the pediatric sub-sample; also includes studies with adults)

- Gibbons, R. D., Brown, C. H., Hur, K., Davis, J. M., & Mann, J. J. (2012). Suicidal thoughts and behavior with antidepressant treatment: reanalysis of the randomized placebo-controlled studies of fluoxetine and venlafaxine. *Archives of general psychiatry*, 69(6), 580-587.

Ages 8-12

- Davis, N. O., Bower, J., Kollins, S. H. (2018). Proof-of-concept study of an at-home, engaging, digital intervention for pediatric ADHD. *PLoS ONE*, 13(1).
- Soffer SL, Lewis J, Lawrence OS, Marroquin YA, Doupnik SK, Benton TD. (2022) Assessing Suicide Risk in a Pediatric Outpatient Behavioral Health System: A Quality Improvement Report. *Pediatr Qual Saf*. Jun 14;7(3) **[mean age 10.6 years; median 14.7 years; range 8.6 to 20.8]**
- Flamarique, I., Santosh, P., Zuddas, A. *et al.* Development and psychometric properties of the Suicidality: Treatment Occurring in Paediatrics (STOP) Suicidality Assessment Scale (STOP-SAS) in children and adolescents. *BMC Pediatr* **16**, 213 (2016). [8-11 group and 12-17 group]

Ages 10-18

- Scott, M., Underwood, M., & Lamis, D. A. (2015). Suicide and Related-Behavior Among Youth Involved in the Juvenile Justice System. *Child and Adolescent Social Work Journal*, 32(6), 517-527.
- Jakobsen, I. S., Larsen, K. J., & Horwood, J. L. (2017). Suicide Risk Assessment in Adolescents—C-SSRS, K10, and READ. *Crisis*.
- Schwartzman, J. M., Muscatello, R. A., & Corbett, B. A. (2023). Assessing suicidal thoughts and behaviors and nonsuicidal self-injury in autistic and non-autistic early adolescents using the Columbia Suicide Severity Rating Scale. *Autism [10-13 year olds]*

Ages 11-17

- Goodyer, I. M., Tsancheva, S., Byford, S., Dubicka, B., Hill, J., Kelvin, R., ... & Wilkinson, P. (2011). Improving mood with psychoanalytic and cognitive therapies (IMPACT): a pragmatic effectiveness superiority trial to investigate whether specialised psychological treatment reduces the risk for relapse in adolescents with moderate to severe unipolar depression: study protocol for a randomised controlled trial. *Trials*, 12(1), 175.

Ages 12-17

- Crandal, B. R., Aguinaldo, L. D., Carter, C., Billman, G. F., Sanderson, K., & Kuelbs, C. (2022). Opportunities for Early Identification: Implementing Universal Depression Screening with a Pathway to Suicide Risk Screening in a Pediatric Health Care System. *The Journal of Pediatrics*, 241, 29-35.

- Findling, R. L., Cutler, A. J., Saylor, K., Gasior, M., Hamdani, M., Ferreira-Cornwell, M. C., & Childress, A. C. (2013). A long-term open-label safety and effectiveness trial of lisdexamfetamine dimesylate in adolescents with attention-deficit/hyperactivity disorder. *J of child and adolescent psychopharmacology*, 23(1), 11-21.
- Findling, R.L., A. Robb, and A. Bose, *Escitalopram in the treatment of adolescent depression: a randomized, double-blind, placebo-controlled extension trial*. *J Child Adolesc Psychopharmacol*, 2013. 23(7): p. 468-80.
- Emslie, G. J., Ventura, D., Korotzer, A., & Tourkodimitris, S. (2009). Escitalopram in the treatment of adolescent depression: a randomized placebo-controlled multisite trial. *Journal of the American Academy of Child & Adolescent Psychiatry*, 48(7), 721-729.

Ages 12-17.5

- Glennon, J., Purper-Ouakil, D., Bakker, M., Zuddas, A., Hoekstra, P., Schulze, U., ... & Coghill, D. (2014). Paediatric European Risperidone Studies (PERS): context, rationale, objectives, strategy, and challenges. *European child & adolescent psychiatry*, 23(12), 1149-1160.

Ages 12-18

- Posner, K., Brown, G. K., Stanley, B., Brent, D. A., Yershova, K. V., Oquendo, M. A., ... & Mann, J. J. (2011). The Columbia–Suicide Severity Rating Scale: initial validity and internal consistency findings from three multisite studies with adolescents and adults. *American Journal of Psychiatry*, 168(12), 1266-1277.
- Brent, D., Emslie, G., Clarke, G., Asarnow, J., Spirito, A., Ritz, L., ... & Keller, M. (2009). Predictors of spontaneous and systematically assessed suicidal adverse events in the treatment of SSRI-resistant depression in adolescents (TORDIA) study. *American Journal of Psychiatry*, 166(4), 418-426.
- Brent, D. A., Greenhill, L. L., Compton, S., Emslie, G., Wells, K., Walkup, J. T., ... & Turner, J. B. (2009). The Treatment of Adolescent Suicide Attempters study (TASA): predictors of suicidal events in an open treatment trial. *Journal of the American Academy of Child & Adolescent Psychiatry*, 48(10), 987-996.
- Patel A, Watts C, Shiddell S, Couch K, Smith AM, Moran MJ, Connors GP. Universal Adolescent Suicide Screening in a Pediatric Urgent Care Center. *Arch Suicide Res*. 2018 Jan-Mar;22(1):118-127 [ages 12-19]
- Gunes A, Kilincaslan A, Eskin M (2015). Psychometric Properties of the Turkish Version of Columbia-Suicide Severity Rating Scale Among 12-18 year-old adolescents in Turkey. AACAP 62nd Annual Meeting, San Antonio, TX.

Ages 13-17

- King, C. A., Jiang, Q., Czyz, E. K., & Kerr, D. C. (2014). Suicidal ideation of psychiatrically hospitalized adolescents has one-year predictive validity for suicide attempts in girls only. *Journal of abnormal child psychology*, 42(3), 467-477.
- Gipson, P. Y., Agarwala, P., Opperman, K. J., Horwitz, A., & King, C. A. (2014). Columbia-Suicide Severity Rating Scale: Predictive Validity with Adolescent Psychiatric Emergency Patients. *Pediatric emergency care*, 31:2, 88-93.
- Rabinovitch, S. M., Kerr, D. C., Leve, L. D., & Chamberlain, P. (2014). Suicidal behavior outcomes of childhood sexual abuse: Longitudinal study of adjudicated girls. *Suicide and life-threatening behavior*.
- Pestian, J. P., Grupp-Phelan, J., Bretonnel Cohen, K., Meyers, G., Richey, L. A., Matykiewicz, P., & Sorter, M. T. (2016). A controlled trial using natural language processing to examine the language of suicidal adolescents in the emergency department. *Suicide and life-threatening behavior*, 46(2), 154-159.

Ages 13-19

- Croarkin, P. E., Nakonezny, P. A., Deng, Z. D., Romanowicz, M., Voort, J. L. V., Camsari, D. D., ... & Lewis, C. P. (2018). High-frequency repetitive TMS for suicidal ideation in adolescents with depression. *Journal of affective disorders*, 239, 282-290.

Ages 14-19

- Kondo, D. G., Sung, Y. H., Hellem, T. L., Fiedler, K. K., Shi, X., Jeong, E. K., & Renshaw, P. F. (2011). Open-label adjunctive creatine for female adolescents with SSRI-resistant major depressive disorder: a 31-phosphorus magnetic resonance spectroscopy study. *Journal of affective disorders*, 135(1), 354-361.
- Kaplow, J. B., Gipson, P. Y., Horwitz, A. G., Burch, B. N., & King, C. A. (2014). Emotional suppression mediates the relation between adverse life events and adolescent suicide: Implications for prevention. *Prevention Science*, 15(2), 177-185.
- King, C. A., Berona, J., Czyz, E., Horwitz, A. G., & Gipson, P. Y. (2015). Identifying Adolescents at Highly Elevated Risk for Suicidal Behavior in the Emergency Department. *Journal of child and adolescent psychopharmacology*.
- Hooper Weatherly, A. (2019). Effectiveness of Two Psychiatric Screening Tools for Adolescent Suicide Risk. *Pediatric Nursing*, 45(4).

Studies with Adolescents and Young Adults

Ages 11-21

- Hill et al (2021) Suicide ideation and attempts in a pediatric emergency department before and during COVID-19. *Pediatrics*, 147.
- Hill et al (2021). Geospatial identification of high youth suicide risk areas via electronic health records: avenues for research and prevention efforts. *Suicide Life Threaten Behav.*, 51:255–62.

Ages 13-25

- Czyz, E. K., Horwitz, A. G., Arango, A., Cole-Lewis, Y., Berona, J., & King, C. A. (2016). Coping with suicidal urges among youth seen in a psychiatric emergency department. *Psychiatry research*, 241, 175-181.

Ages 14-39

- Lucas, M. S., Brawner, B. M., Hardie, T. L., Beacham, B., Paidipati, C., Diaz, M., ... & Deatrck, J. A. (2015, September). Assessing Suicidal Ideation and Behaviors Among Survivors of Childhood Brain Tumors and Their Mothers During Sociobehavioral Research. In *Oncology nursing forum* (Vol. 42, No. 5, pp. E319-29).

Ages 15-20

- Gray, K. M., Carpenter, M. J., Lewis, A. L., Klintworth, E. M., & Upadhyaya, H. P. (2012). Varenicline versus bupropion XL for smoking cessation in older adolescents: A randomized, double-blind pilot trial. *Nicotine & Tobacco Research*, 14(2), 234-239.

Ages 15-24

- Horwitz, A. G., Czyz, E. K., & King, C. A. (2015). Predicting future suicide attempts among adolescent and emerging adult psychiatric emergency patients. *Journal of Clinical Child & Adolescent Psychology*, 44:5, 751-761.

Ages 20-22

- Cáceda, R., Durand, D., Cortes, E., Prendes-Alvarez, S., Moskovciak, T., Harvey, P. D., & Nemeroff, C. B. (2014). Impulsive choice and psychological pain in acutely suicidal depressed patients. *Psychosomatic medicine*, 76(6), 445-451.

Demographic Groups

Asian Americans

- Maru, M., Saraiya, T., Lee, C.S., Meghani, O., Hien, D. and Hahm, H.C., 2018. The relationship between intimate partner violence and suicidal ideation among young Chinese, Korean, and Vietnamese American women. *Women & Therapy*, pp.1-17.

School Protocols

- Stargell, N. A, Zoldan C. A., Kress, V. E., Walker-Andrews, M., and Whisenhunt, J. L. (2017). Student Non-Suicidal Self-Injury: A Protocol for School Counselors (Conceptual). *Professional School Counseling*, 21(1), 37-46.
- Rhode Island?

Medical Specialties

Neurology

- Hesdorffer, D. C., French, J. A., Posner, K., DiVentura, B., Pollard, J. R., Sperling, M. R., & Kanner, A. M. (2013). "Suicidal ideation and behavior screening in intractable focal **epilepsy** eligible for drug trials." *Epilepsia*, 1-9.
- Pereira, A., Gitlin, M. J., Gross, R. A., Posner, K., & Dworkin, R. H. (2013). "Suicidality associated with antiepileptic drugs: Implications for the treatment of **neuropathic pain and fibromyalgia**." *PAIN*[®], 154(3), 345-349.
- Skljarevski, V., Zhang, S., Desai, D., Alaka, K. J., Palacios, S., Miazgowski, T., & Patrick, K. (2010). Duloxetine versus placebo in patients with **chronic low back pain**: a 12-week, fixed-dose, randomized, double-blind trial. *The Journal of Pain*, 11(12), 1282-1290.
- Legarreta, M., Bueler, E., DiMuzio, J., McGlade, E. and Yurgelun-Todd, D., 2018. Suicide Behavior and Chronic Pain: An Exploration of Pain-Related Catastrophic Thinking, Disability, and Descriptions of the Pain Experience. *The Journal of nervous and mental disease*, 206(3), pp.217-222.
- Dale, M., & van Duijn, E. (2015). Anxiety in **Huntington's** disease. *The Journal of neuropsychiatry and clinical neurosciences*, 27(4), 262-271.
- van Duijn, E., Vrijmoeth, E. M., Giltay, E. J., Landwehrmeyer, G. B., & Registry investigators of the European **Huntington's** Disease Network. (2018). Suicidal ideation and suicidal behavior according to the C-SSRS in a European cohort of Huntington's disease gene expansion carriers. *Journal of affective disorders*, 228, 194-204.
- Elfil, M., Ahmed, N., Alapati, A., Bahekar, R., Kandil, M., Kim, C., ... & Louis, E. D. (2019). Suicidal risk and demoralization in **Parkinson disease**. *Journal of neurology*, 1-9.

Oncology

- Lucas, M. S., Brawner, B. M., Hardie, T. L., Beacham, B., Paidipati, C., Diaz, M., ... & Deatrck, J. A. (2015, September). Assessing Suicidal Ideation and Behaviors Among Survivors of **Childhood Brain Tumors** and Their Mothers During Sociobehavioral Research. In *Oncology nursing forum* (Vol. 42, No. 5, pp. E319-29).

Psoriasis

- Pompili, M., Innamorati, M., Forte, A., Erbuto, D., Lamis, D. A., Narcisi, A., ... & Bellini, S. (2017). Psychiatric comorbidity and suicidal ideation in psoriasis, melanoma and allergic disorders. *International journal of psychiatry in clinical practice*, 21(3), 209-214.

Psychiatric Conditions

Alzheimer's

- Nave, S., Doody, R. S., Boada, M., Grimmer, T., Savola, J. M., Delmar, P., ... & Ricci, B. (2017). Sembragiline in Moderate Alzheimer's Disease: Results of a Randomized, Double-Blind, Placebo-Controlled Phase II Trial (MAyflOwer RoAD). *Journal of Alzheimer's Disease*, (Preprint), 1-12.
- Delnomdedieu, M., Duvvuri, S., Li, D. J., Atassi, N., Lu, M., Brashear, H. R., ... & Kupiec, J. W. (2016). First-In-Human safety and long-term exposure data for AAB-003 (PF-05236812) and biomarkers after intravenous infusions of escalating doses in patients with mild to moderate Alzheimer's disease. *Alzheimer's research & therapy*, 8(1), 12.
- Kim, S. Y., Choi, S. H., Rollema, H., Schwam, E. M., McRae, T., Dubrava, S., & Jacobsen, J. (2013). Phase II crossover trial of varenicline in mild-to-moderate **Alzheimer's** disease. *Dementia and geriatric cognitive disorders*, 37(3-4), 232-245.

ADHD

- Wigal, S. B., Nordbrock, E., Adjei, A. L., Childress, A., Kupper, R. J., & Greenhill, L. (2015). Efficacy of methylphenidate hydrochloride extended-release capsules (Aptensio XR™) in **children and adolescents with attention-deficit/hyperactivity disorder**: A phase III, randomized, double-blind study. *CNS drugs*, 29(4), 331-340.
- Sangal, R. B., Blumer, J. L., Lankford, D. A., Grinnell, T. A., & Huang, H. (2014). Eszopiclone for insomnia associated with **attention-deficit/hyperactivity disorder**. *Pediatrics*, 134(4), e1095-e1103.
- Davis, N. O., Bower, J., Kollins, S. H. (2018). Proof-of-concept study of an at-home, engaging, digital intervention for pediatric ADHD. *PLoS ONE*, 13(1).

Alcohol

- Khemiri, L., Jokinen, J., Runeson, B., & Jayaram-Lindström, N. (2016). Suicide risk associated with experience of violence and impulsivity in alcohol dependent patients. *Scientific reports*, 6, 19373.

Autism

- Danforth, A. L., Struble, C. M., Yazar-Klosinski, B., & Grob, C. S. (2015). MDMA-assisted therapy: A new treatment model for social anxiety in autistic adults. *Progress in Neuro-Psychopharmacology and Biological Psychiatry*.
- Schwartzman, J. M., Muscatello, R. A., & Corbett, B. A. (2023). Assessing suicidal thoughts and behaviors and nonsuicidal self-injury in autistic and non-autistic early adolescents using the Columbia Suicide Severity Rating Scale. *Autism*

Bipolar Depression

- Harvey, P. D., Siever, L. J., Huang, G. D., Muralidhar, S., Zhao, H., Miller, P., ... & Brophy, M. (2014). The genetics of functional disability in schizophrenia and bipolar illness: methods and initial results for VA cooperative study# 572. *American Journal of Medical Genetics Part B: Neuropsychiatric Genetics*, 165(4), 381-389.
- Earley, W., Durgam, S., Lu, K., Debelle, M., Laszlovszky, I., Vieta, E., & Yatham, L. N. (2017). Tolerability of Cariprazine in the Treatment of Acute Bipolar I Mania: A Pooled Post Hoc Analysis of 3 Phase II/III Studies. *Journal of Affective Disorders*.

- Weinstein, S. M., Henry, D. B., Katz, A. C., Peters, A. T., & West, A. E. (2015). Treatment moderators of child-and family-focused cognitive-behavioral therapy for **pediatric bipolar disorder**. *Journal of the American Academy of Child & Adolescent Psychiatry*, 54(2), 116-125.

Complicated Grief

- Supiano, K. P., & Luptak, M. (2013). Complicated grief in older adults: A randomized controlled trial of complicated grief group therapy. *The Gerontologist*, 54(5), 840-856. (6-item screener version)

Generalized Anxiety Disorder (GAD)

- Strawn, J. R., Prakash, A., Zhang, Q., Pangallo, B. A., Stroud, C. E., Cai, N., & Findling, R. L. (2015). A randomized, placebo-controlled study of duloxetine for the treatment of **children and adolescents with generalized anxiety disorder**. *Journal of the American Academy of Child & Adolescent Psychiatry*, 54(4), 283-293.

Perinatal/Postpartum Depression

- Kaner, S. J., Colquhoun, H., Doherty, J., Raines, S., Hoffmann, E., Rubinow, D. R., & Meltzer-Brody, S. (2017). Open-label, proof-of-concept study of brexanolone in the treatment of severe postpartum depression. *Human Psychopharmacology: Clinical and Experimental*, 32(2).
- Kim, J. J., Silver, R. K., Elue, R., Adams, M. G., La Porte, L. M., Cai, L., ... & Gibbons, R. D. (2016). The experience of depression, anxiety, and mania among perinatal women. *Archives of women's mental health*, 19(5), 883-890.
- Ghahramanlou-Holloway, M., Cox, D. W., Fritz, E. C., & George, B. J. (2011). An evidence-informed guide for working with military women and veterans. *Professional Psychology: Research and Practice*, 42(1), 1.

Psychosis/Schizophrenia

- DeVylder, J. E., Jahn, D. R., Doherty, T., Wilson, C. S., Wilcox, H. C., Schiffman, J., & Hilimire, M. R. (2015). Social and psychological contributions to the co-occurrence of sub-threshold psychotic experiences and suicidal behavior. *Social psychiatry and psychiatric epidemiology*, 1-12. [young adult SELF-REPORT CSSRS]
- Harvey, P. D., Siever, L. J., Huang, G. D., Muralidhar, S., Zhao, H., Miller, P., ... & Brophy, M. (2014). The genetics of functional disability in schizophrenia and bipolar illness: methods and initial results for VA cooperative study# 572. *American Journal of Medical Genetics Part B: Neuropsychiatric Genetics*, 165(4), 381-389.
- Hettige, N. C., Bani-Fatemi, A., Kennedy, J. L., & De Luca, V. (2017). Assessing the risk for suicide in schizophrenia according to migration, ethnicity and geographical ancestry. *BMC Psychiatry*, 17(1), 63.

PTSD

- Legarreta, M., Graham, J., North, L., Bueler, C. E., McGlade, E., & Yurgelun-Todd, D. (2015). DSM-5 posttraumatic stress disorder symptoms associated with suicide behaviors in veterans. *Psychological trauma: theory, research, practice, and policy*, 7(3), 277.

Sleep

- Ellis, T. E., Rufino, K. A., & Nadorff, M. R. (2017). Treatment of Nightmares in Psychiatric Inpatients With Imagery Rehearsal Therapy: An Open Trial and Case Series. *Behavioral Sleep Medicine*, 1-14.
- Sangal, R. B., Blumer, J. L., Lankford, D. A., Grinnell, T. A., & Huang, H. (2014). Eszopiclone for insomnia associated with **attention-deficit/hyperactivity disorder**. *Pediatrics*, 134(4), e1095-e1103.

Healthcare Systems

- Rossom, R. C., Simon, G. E., Beck, A., Ahmedani, B. K., Steinfeld, B., Trangle, M., & Solberg, L. (2016). Facilitating Action for Suicide Prevention by Learning Health Care Systems. *Psychiatric Services*.

Outpatient Settings

Outpatient Psychiatry

- Viguera, A. C., Milano, N., Laurel, R., Thompson, N. R., Griffith, S. D., Baldessarini, R. J., & Katzan, I. L. (2015). Comparison of electronic screening for suicidal risk with the Patient Health Questionnaire Item 9 and the Columbia Suicide Severity Rating Scale in an outpatient psychiatric clinic. *Psychosomatics*, 56(5), 460-469.

Integrated Primary Care

- Kearney, L. K., Wray, L. O., Dollar, K. M., & King, P. R. (2015). Establishing Measurement-based Care in Integrated Primary Care: Monitoring Clinical Outcomes Over Time. *Journal of clinical psychology in medical settings*, 22(4), 213-227.

Veterans

- Zisook, S., Tal, I., Weingart, K., Hicks, P., Davis, L. L., Chen, P., ... & Pilkinton, P. D. (2016). Characteristics of US Veteran Patients with Major Depressive Disorder who require “next-step” treatments: A VAST-D report. *Journal of Affective Disorders*, 206, 232-240.
- Legarreta, M., Graham, J., North, L., Bueler, C. E., McGlade, E., & Yurgelun-Todd, D. (2015). DSM–5 posttraumatic stress disorder symptoms associated with suicide behaviors in veterans. *Psychological trauma: theory, research, practice, and policy*, 7(3), 277.
- Harvey, P. D., Siever, L. J., Huang, G. D., Muralidhar, S., Zhao, H., Miller, P., ... & Brophy, M. (2014). The genetics of functional disability in schizophrenia and bipolar illness: methods and initial results for VA cooperative study# 572. *American Journal of Medical Genetics Part B: Neuropsychiatric Genetics*, 165(4), 381-389.
- Legarreta, M., Graham, J., North, L., Bueler, C. E., McGlade, E., & Yurgelun-Todd, D. (2015). DSM–5 posttraumatic stress disorder symptoms associated with suicide behaviors in veterans. *Psychological trauma: theory, research, practice, and policy*, 7(3), 277.
- Goodman, M., Banthin, D., Blair, N. J., Mascitelli, K. A., Wilsnack, J., Chen, J., ... & New, A. S. (2016). A randomized trial of dialectical behavior therapy in high-risk suicidal veterans. *The Journal of Clinical Psychiatry*, 77(12), 4031.
- Kimbrel, N. A., DeBeer, B. B., Meyer, E. C., Gulliver, S. B., & Morissette, S. B. (2016). Nonsuicidal self-injury and suicide attempts in Iraq/Afghanistan war veterans. *Psychiatry research*, 243, 232-237.
- Reist, C., Mee, S., Fujimoto, K., Rajani, V., Bunney, W. E., & Bunney, B. G. (2017). Assessment of psychological pain in suicidal veterans. *PLoS one*, 12(5), e0177974.
- Pigeon, W. R., Funderburk, J., Bishop, T. M., & Crean, H. F. (2017). Brief cognitive behavioral therapy for insomnia delivered to depressed veterans receiving primary care services: a pilot study. *Journal of affective disorders*, 217, 105-111.
- DeBeer, B. B., Meyer, E. C., Kimbrel, N. A., Kittel, J. A., Gulliver, S. B., & Morissette, S. B. (2018). Psychological inflexibility predicts of suicidal ideation over time in veterans of the conflicts in Iraq and Afghanistan. *Suicide and Life-threatening Behavior*, 48(6), 627-641.
- Sheth, C., Prescott, A., Bueler, E., DiMuzio, J., Legarreta, M., Renshaw, P. F., ... & McGlade, E. (2018). Alterations in anterior cingulate cortex myoinositol and aggression in veterans with suicidal behavior: a proton magnetic resonance spectroscopy study. *Psychiatry Research: Neuroimaging*, 276, 24-32.

- Yesavage, J. A., Fairchild, J. K., Mi, Z., Biswas, K., Davis-Karim, A., Phibbs, C. S., ... & VA Cooperative Studies Program Study Team. (2018). Effect of repetitive transcranial magnetic stimulation on treatment-resistant major depression in US veterans: a randomized clinical trial. *JAMA psychiatry*, 75(9), 884-893.
- Hwang, J., Legarreta, M., Bueler, C. E., DiMuzio, J., McGlade, E., Lyoo, I. K., & Yurgelun-Todd, D. (2018). Increased efficiency of brain connectivity networks in veterans with suicide attempts. *Neuroimage: clinical*, 20, 318-326.
- Kachadourian, L. K., Gandelman, E., Ralevski, E., & Petrakis, I. L. (2018). Suicidal ideation in military veterans with alcohol dependence and PTSD: The role of hostility. *The American journal on addictions*, 27(2), 124-130.
- Kopacz, M. S., Crean, H. F., L. Park, C., & Hoff, R. A. (2018). Religious coping and suicide risk in a sample of recently returned veterans. *Archives of suicide research*, 22(4), 615-627.
- Kimbrel, N. A., Meyer, E. C., DeBeer, B. B., Gulliver, S. B., & Morissette, S. B. (2018). The impact of cannabis use disorder on suicidal and nonsuicidal self-injury in Iraq/Afghanistan-era veterans with and without mental health disorders. *Suicide and Life-Threatening Behavior*, 48(2), 140-148.
- Harvey, P. D., Posner, K., Rajeevan, N., Yershova, K. V., Aslan, M., & Concato, J. (2018). Suicidal ideation and behavior in US veterans with schizophrenia or bipolar disorder. *Journal of psychiatric research*, 102, 216-222.
- Matarazzo, B. B., Brown, G. K., Stanley, B., Forster, J. E., Billera, M., Currier, G. W., ... & Brenner, L. A. (2019). Predictive validity of the columbia-suicide severity rating scale among a cohort of at-risk veterans. *Suicide and Life-Threatening Behavior*, 49(5), 1255-1265.
- Hoffmire, C. A., Monteith, L. L., Holliday, R., Park, C. L., Brenner, L. A., & Hoff, R. A. (2019). Administrative Military Discharge and Suicidal Ideation Among Post-9/11 Veterans. *American journal of preventive medicine*, 56(5), 727-735.
- Law, K. C., Allan, N. P., Kolnogorova, K., & Stecker, T. (2019). An examination of PTSD symptoms and their effects on suicidal ideation and behavior in non-treatment seeking veterans. *Psychiatry research*, 274, 12-19.
- Chesin, M., Interian, A., Kline, A., Hill, L. S., King, A., Miller, R., ... & Stanley, B. (2019). Past-year opioid misuse and suicide attempt are positively associated in high suicide risk veterans who endorse past-year substance use. *Addictive behaviors*, 99, 106064.
- Bishop, T. M., Crean, H. F., Hoff, R. A., & Pigeon, W. R. (2019). Suicidal ideation among recently returned veterans and its relationship to insomnia and depression. *Psychiatry research*, 276, 250-261.
- Katz, I., Barry, C. N., Cooper, S. A., Kaspro, W. J., & Hoff, R. A. (2020). Use of the Columbia-Suicide Severity Rating Scale (C-SSRS) in a large sample of Veterans receiving mental health services in the Veterans Health Administration. *Suicide and Life-Threatening Behavior*, 50(1), 111-121.
- Aslan, M., Radhakrishnan, K., Rajeevan, N., Sueiro, M., Goulet, J. L., Li, Y., ... & Harvey, P. D. (2020). Suicidal ideation, behavior, and mortality in male and female US veterans with severe mental illness. *Journal of affective disorders*, 267, 144-152.
- Szpunar, M. J., Crawford, J. N., Baca, S. A., & Lang, A. J. (2020). Suicidal ideation in pregnant and postpartum women veterans: an initial clinical needs assessment. *Military medicine*, 185(1-2), e105-e111.
- Simon, N. M., Hoepfner, S. S., Lubin, R. E., Robinaugh, D. J., Malgaroli, M., Norman, S. B., ... & Rauch, S. A. (2020). Understanding the impact of complicated grief on combat related posttraumatic stress disorder, guilt, suicide, and functional impairment in a clinical trial of post-9/11 service members and veterans. *Depression and anxiety*, 37(1), 63-72.
- Hoffmire, C. A., Monteith, L. L., Forster, J. E., Bernhard, P. A., Blosnich, J. R., Vogt, D., ... & Schneiderman, A. I. (2021). Gender differences in lifetime prevalence and onset timing of suicidal ideation and suicide attempt among post-9/11 veterans and nonveterans. *Medical care*, 59, S84-S91.
- Blosnich, J. R., Garfin, D. R., Maguen, S., Vogt, D., Dichter, M. E., Hoffmire, C. A., ... & Schneiderman, A. (2021). Differences in childhood adversity, suicidal ideation, and suicide attempt among veterans and nonveterans. *American Psychologist*, 76(2), 284.

Forensic Settings

- SEE ALSO “JUVENILE JUSTICE”
- Wilson, N. J. (2017). The Columbia-Suicide Severity Rating Scale: Validation for use as a screen for suicide risk in New Zealand prisons and probation settings. *Practice, 5*(2).

Juvenile Justice

- Kerr, D. C., Gibson, B., Leve, L. D., & DeGarmo, D. S. (2014). Young adult follow-up of adolescent girls in juvenile justice using the Columbia Suicide Severity Rating Scale. *Suicide and life-threatening behavior, 44*(2), 113-129.
- Kerr, D. C., DeGarmo, D. S., Leve, L. D., & Chamberlain, P. (2014). Juvenile justice girls' depressive symptoms and suicidal ideation 9 years after multidimensional treatment foster care. *Journal of consulting and clinical psychology, 82*(4), 684.
- Rabinovitch, S. M., Kerr, D. C., Leve, L. D., & Chamberlain, P. (2014). Suicidal behavior outcomes of childhood sexual abuse: Longitudinal study of adjudicated girls. *Suicide and life-threatening behavior.*
- Scott, M., Underwood, M., & Lamis, D. A. (2015). Suicide and Related-Behavior Among Youth Involved in the Juvenile Justice System. *Child and Adolescent Social Work Journal, 32*(6), 517-527.

Emergency Departments

- Patel, A., Watts, C., Shiddell, S., Couch, K., Smith, A. M., Moran, M. J., & Conners, G. P. (2017). Universal adolescent suicide screening in a pediatric urgent care center. *Archives of suicide research, 1*-10.
- Pestian, J. P., Grupp-Phelan, J., Bretonnel Cohen, K., Meyers, G., Richey, L. A., Matykiewicz, P., & Sorter, M. T. (2016). A controlled trial using natural language processing to examine the language of suicidal adolescents in the emergency department. *Suicide and life-threatening behavior, 46*(2), 154-159.
- Czyz, E. K., Horwitz, A. G., Arango, A., Cole-Lewis, Y., Berona, J., & King, C. A. (2016). Coping with suicidal urges among youth seen in a psychiatric emergency department. *Psychiatry research, 241*, 175-181.
- Arias, S. A., Miller, I., Camargo Jr, C. A., Sullivan, A. F., Goldstein, A. B., Allen, M. H., ... & Boudreaux, E. D. (2015). Factors Associated with Suicide Outcomes 12 Months After Screening Positive for Suicide Risk in the Emergency Department. *Psychiatric Services.*
- Brown, G. K., Currier, G. W., Jager-Hyman, S., & Stanley, B. (2015). Detection and classification of suicidal behavior and nonsuicidal self-injury behavior in emergency departments. *The Journal of clinical psychiatry, 76*(10), 1-478.
- Horwitz, A. G., Czyz, E. K., & King, C. A. (2015). Predicting future suicide attempts among adolescent and emerging adult psychiatric emergency patients. *Journal of Clinical Child & Adolescent Psychology, 44*(5), 751-761.
- King, C. A., Berona, J., Czyz, E., Horwitz, A. G., & Gipson, P. Y. (2015). Identifying adolescents at highly elevated risk for suicidal behavior in the emergency department. *Journal of child and adolescent psychopharmacology, 25*(2), 100-108.
- Gipson, P. Y., Agarwala, P., Opperman, K. J., Horwitz, A., & King, C. A. (2015). Columbia-suicide severity rating scale: predictive validity with adolescent psychiatric emergency patients. *Pediatric emergency care, 31*(2), 88-94.
- Lim, M., Kim, S. W., Nam, Y. Y., Moon, E., Yu, J., Lee, S., ... & Ahn, Y. M. (2014). Reasons for desiring death: examining causative factors of suicide attempters treated in emergency rooms in Korea. *Journal of affective disorders, 168*, 349-356.
- Ducasse, D., René, E., Béziat, S., Guillaume, S., Courtet, P., & Olié, E. (2014). Acceptance and commitment therapy for management of suicidal patients: a pilot study. *Psychotherapy and psychosomatics, 83*(6), 374-376.

- Boudreaux, E. D., Miller, I., Goldstein, A. B., Sullivan, A. F., Allen, M. H., Manton, A. P., ... & Camargo, C. A. (2013). The emergency department safety assessment and follow-up evaluation (ED-SAFE): method and design considerations. *Contemporary clinical trials*, 36(1), 14-24.
- Posner, K., Brown, G. K., Stanley, B., Brent, D. A., Yershova, K. V., Oquendo, M. A., ... & Mann, J. J. (2011). The Columbia–Suicide Severity Rating Scale: initial validity and internal consistency findings from three multisite studies with adolescents and adults. *American Journal of Psychiatry*, 168(12), 1266-1277.

In-Patient Psychiatric Settings

- Allen, J. G., Fowler, J. C., Madan, A., Ellis, T. E., Oldham, J. M., & Frueh, B. C. (2017). Discovering the impact of psychotherapeutic hospital treatment for adults with serious mental illness. *Bulletin of the Menninger Clinic*, 81(1), 1-38.
- Madan, A., Frueh, B. C., Allen, J. G., Ellis, T. E., Rufino, K. A., Oldham, J. M., & Fowler, J. C. (2016). Psychometric reevaluation of the Columbia-Suicide Severity Rating Scale: findings from a prospective, inpatient cohort of severely mentally ill adults. *Journal of clinical psychiatry*, 77(7), e867-e873.
- Lucas, M. S., Brawner, B. M., Hardie, T. L., Beacham, B., Paidipati, C., Diaz, M., ... & Deatrck, J. A. (2015, September). Assessing Suicidal Ideation and Behaviors Among Survivors of Childhood Brain Tumors and Their Mothers During Sociobehavioral Research. In *Oncology nursing forum* (Vol. 42, No. 5, pp. E319-29).
- Teti, G. L., Rebok, F., Grndas, L. N., Rodante, D., Fogola, A., & Daray, F. M. (2014). Patients hospitalized for suicidal ideation and suicide attempt in a Mental Health Hospital: Clinico-demographical features and 6-month follow-up. *Vertex*, 25(115), 203-212.
- Arias, S. A., Zhang, Z., Hillerns, C., Sullivan, A. F., Boudreaux, E. D., Miller, I., & Camargo, C. A. (2014). Using structured telephone follow-up assessments to improve suicide-related adverse event detection. *Suicide and life-threatening behavior*, 44(5), 537-547.
- King, C. A., Jiang, Q., Cxyz, E. K., & Kerr, D. C. (2014). Suicidal ideation of psychiatrically hospitalized adolescents has one-year predictive validity for suicide attempts in girls only. *Journal of abnormal child psychology*, 42(3), 467-477.
- Cáceda, R., Durand, D., Cortes, E., Prendes-Alvarez, S., Moskovciak, T., Harvey, P. D., & Nemeroff, C. B. (2014). Impulsive choice and psychological pain in acutely suicidal depressed patients. *Psychosomatic medicine*, 76(6), 445-451.
- Ellis, T. E., Rufino, K. A., Allen, J. G., Fowler, J. C., & Jobes, D. A. (2015). Impact of a suicide-specific intervention within inpatient psychiatric care: the Collaborative Assessment and Management of Suicidality. *Suicide and life-threatening behavior*, 45(5), 556-566.

Mobile Crisis

- Muehsam, J. P. (2018). Association between clinical observations and a mobile crisis team's level of care recommendations. *Community Mental Health Journal*, 1-7.

Telemedicine

- Arias SA, Zhang Z, Hillerns C, Sullivan AF, Boudreaux ED, Miller I, Camargo CA (2014). Using Structured Telephone Follow-up Assessments to Improve Suicide-Related Adverse Event Detection. *Suicide and Life-Threatening Behavior*44(5): 537-47.

Treatment Efficacy for Suicidal Outcomes

- Ionescu, D. F., Swee, M. B., Pavone, K. J., Taylor, N., Akeju, O., Baer, L., ... & Brown, E. N. (2016). Rapid and sustained reductions in current suicidal ideation following repeated doses of intravenous ketamine: secondary analysis of an open-label study. *The Journal of clinical psychiatry*.
- Prakash, A., Lobo, E., Kratochvil, C. J., Tamura, R. N., Pangallo, B. A., Bullok, K. E., Quinlan, T., Emslie, G.J. & March, J. S. (2012). An open-label safety and pharmacokinetics study of duloxetine in pediatric patients with major depression. *Journal of Child and Adolescent Psychopharmacology*, 22(1), 48-55.

- Croarkin, P. E., Nakonezny, P. A., Deng, Z. D., Romanowicz, M., Voort, J. L. V., Camsari, D. D., ... & Lewis, C. P. (2018). High-frequency repetitive TMS for suicidal ideation in adolescents with depression. *Journal of affective disorders*, 239, 282-290.

Regulatory Guidelines for Treatment & Assessment of Suicidal Outcomes

- Wasserman, D., Rihmer, Z., Rujescu, D., Sarchiapone, M., Sokolowski, M., Titelman, D., ... & Carli, V. (2012). The European Psychiatric Association (EPA) guidance on suicide treatment and prevention. *European Psychiatry*, 27(2), 129-141.
- US Food and Drug Administration. *Suicidal Ideation and Behavior: Prospective Assessment of Occurrence in Clinical Trials*. <http://www.fda.gov/Drugs/GuidanceComplianceRegulatoryInformation/Guidances/ucm315156.htm>. Rockville, MD: US Department of Health and Human Services; 2012.

Reviews of Suicide Risk Assessment Tools

- Bolton, J. M., Gunnell, D., & Turecki, G. (2015). Suicide risk assessment and intervention in people with mental illness. *British Medical Journal*, 351(8034).
- Posner, K., Buchanan, J., Amira, L., Yershova, K., Lesser, A., Goldstein, E. (2014). Identification and screening of suicide risk. In S. Koslow, C., Nemeroff, P., Ruiz, (eds.), *A Concise Guide to Understanding Suicide: Epidemiology, Pathophysiology and Prevention*. Cambridge University Press.

Cross-Cultural Settings

Latin America (Spanish)

- Teti, G. L., Rebok, F., Grndas, L. N., Rodante, D., Fogola, A., & Daray, F. M. (2014). Patients hospitalized for suicidal ideation and suicide attempt in a Mental Health Hospital: Clinico-demographical features and 6-month follow-up. *Vertex*, 25(115), 203-212.

Australia

- van Spijker, B. A., Caelear, A. L., Batterham, P. J., Mackinnon, A. J., Gosling, J. A., Kerkhof, A. J., ... & Christensen, H. (2015). Reducing suicidal thoughts in the Australian general population through web-based self-help: study protocol for a randomized controlled trial. *Trials*, 16(1), 589-589.

China

- Wang, H., Xue, Y., Chen, Y., Zhang, R., Wang, H., Zhang, Y., ... & Tan, Q. (2013). Efficacy of repetitive transcranial magnetic stimulation in the prevention of relapse of depression: study protocol for a randomized controlled trial. *Trials*, 14(1), 338.

Chile

- Núñez, D., Arias, V., Méndez-Bustos, P., & Fresno, A. (2019). Is a brief self-report version of the Columbia severity scale useful for screening suicidal ideation in **Chilean** adolescents? *Comprehensive psychiatry*, 88, 39-48.

Croatia

- Sisek-Šprem, M. (2012). Demographic characteristics of aggressive patients with schizophrenia. *Socijalna psihijatrija*, 40(3), 213-220.

Ethiopia

- Borba, C. P., Fekadu, A., Teferra, S., Bekele, D., Shibre, T., Oppenheim, C. E., ... & Henderson, D. C. (2014). A placebo-controlled trial of folate with B12 in patients with schizophrenia with residual symptoms in Ethiopia using a sequential parallel comparison design. *British Journal of Medicine and Medical Research*, 4(23), 4090.

France

- Welniarz, B., & Saintoyan, F. (2015). Dépression de l'enfant et de l'adolescent: place du traitement médicamenteux et de l'hospitalisation. *Neuropsychiatrie de l'Enfance et de l'Adolescence*.

Germany

- von Klitzing, K. (2008). Depressionen im Kindes und Jugendalter. *Kinder-und Jugendmedizin*, 8(1), 18-23.

Hungary

- Wasserman, D., Rihmer, Z., Rujescu, D., Sarchiapone, M., Sokolowski, D.T., Zalsman, G., ... & Carli, V. (2012). Az Európai Pszichiátriai Szövetség (European Psychiatric Association, EPA) útmutatója az öngyilkosság kezelésére és megelőzésére. *Neuropsychopharmacol Hung*,14(2), 113-136.

Indonesia

- Pratiwi, J., & Undarwati, A. (2014). SUICIDE IDEATION PADA REMAJA DI KOTA SEMARANG. *Developmental and Clinical Psychology*, 3(1).

India

- Pasi, S., Singh, P. K., Pandey, R. K., Dikshit, P. C., Jiloha, R. C., & Rao, V. R. (2015). Evaluation of psychiatric and genetic risk factors among primary relatives of suicide completers in Delhi NCR region, India. *Psychiatry research*, 229(3), 933-939.
- Bansal, K. (2013). Pre- and Post-Psychosocial Factors of Subjects enrolled in Phase-III Oncology Clinical Trials. *Journal of Academia and Industrial Research (JAIR)*, 2(5), 312.

Korea

- Lim, M., Lee, S., & Park, J. I. (2015). Characteristics of Korean Suicide Attempters. *Journal of Korean Neuropsychiatric Association*, 54(2), 209-215.

Lebanon

Zakhour, M., Haddad, C., Sacre, H. *et al.* Suicidal ideation among Lebanese adults: scale validation and correlates. *BMC Psychiatry* **21**, 100 (2021). <https://doi.org/10.1186/s12888-021-03111-7>

Spain

- Teti, G. L., Rebok, F., Grndas, L. N., Rodante, D., Fogola, A., & Daray, F. M. (2014). Patients hospitalized for suicidal ideation and suicide attempt in a Mental Health Hospital: Clinico-demographical features and 6-month follow-up. *Vertex*, 25(115), 203-212.

- Ruiz, V., Torra, M., & Bernardo, M. (2016). Respuesta antipsicótica en trastorno delirante y esquizofrenia: estudio prospectivo de cohortes. *Actas Esp Psiquiatr*, 44(4), 125-35.
- Sáiz, P. A., Rodríguez-Revuelta, J., González-Blanco, L., Burón, P., Al-Halabí, S., Garrido, M., ... & Bobes, J. (2014). Protocolo de estudio de un programa para la prevención de la recurrencia del comportamiento suicida basado en el manejo de casos (PSyMAC). *Revista de Psiquiatría y Salud Mental*, 7(3), 131-138.

Sri Lanka

- Suraweera, C., Hanwella, R., Sivayokan, S., & de Silva, V. (2013). Rating Scales validated for Sri Lankan populations. *Sri Lanka J of Psychiatry*, 4(2), 16-24.